

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical  
(Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

<b>Data Requirement:</b>	PMRA Data Code	{.....}
	EPA DP Barcode	358148
	OECD Data Point	{.....}
	EPA MRID	47560119
	EPA Guideline	850.2200

**Test material:** DPX-MAT28 Technical **Purity:** 92.2%  
**Common name** Aminocyclopyrachlor  
**Chemical name:**  
IUPAC: 6-amino-5-chloro-2-cyclopropylpyrimidine-4-carboxylic acid  
CAS: 6-amino-5-chloro-2-cyclopropyl-4-pyrimidinecarboxylic acid  
CAS No.: 858956-08-8  
Synonyms: None reported

**Primary Reviewer:** Christie E. Padova  
Staff Scientist, Dynamac Corporation

**Signature:** *Christie E. Padova*  
**Date:** 07/10/09

**Primary Reviewer:** Teri S. Myers  
Senior Scientist, Cambridge Environmental Inc.

**Signature:** *Teri S. Myers*  
**Date:** 07/23/09

**Secondary Reviewer:** Anita Ullagaddi  
EPA/OPP/EFED/ERB1

**Signature:** *Anita*  
**Date:** 10/05/09

**Reference/Submission No.:** {.....}

<b>Company Code</b>	{.....}	[For PMRA]
<b>Active Code</b>	{.....}	[For PMRA]
<b>Use Site Category</b>	{.....}	[For PMRA]
<b>EPA PC Code</b>	None	

**Date Evaluation Completed:** 10/05/09

**CITATION:** Hubbard, P.M., K.H. Martin, and J.B. Beavers. 2007. DPX-MAT28 Technical: A Dietary LC<sub>50</sub> Study with the Mallard. Unpublished study performed by Wildlife International Ltd., Easton, MD. Laboratory Project No. 112-598. Study sponsored by E.I. du Pont de Nemours and Company, Wilmington, DE. Study initiated March 28, 2007 and submitted June 6, 2007.

**DISCLAIMER:** This document provides guidance for EPA and PMRA reviewers on how to complete a data evaluation record after reviewing a scientific study concerning the acute dietary toxicity of a pesticide to avian species. It is not intended to prescribe conditions to any external party for conducting this study nor to establish absolute criteria regarding the assessment of whether the study is scientifically sound and whether the study satisfies any applicable data requirements. Reviewers are expected to review and to determine for each study, on a case-by-case basis, whether it is scientifically sound and provides sufficient information to satisfy applicable data requirements. Studies that fail to meet any of the conditions may be accepted, if appropriate; similarly, studies that meet all of the conditions may be rejected, if appropriate. In sum, the reviewer is to take into account the totality of factors related to the test methodology and results in determining the acceptability of the study.



2084708

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical  
(Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

**EXECUTIVE SUMMARY:**

The acute dietary toxicity of DPX-MAT28 Technical (aminocyclopyrachlor) to 10-day old mallard duck (*Anas platyrhynchos*) was assessed over 8 days. DPX-MAT28 was administered to the birds in the diet at nominal concentrations of 0 (negative control), 562, 1000, 1780, 3160, and 5620 mg ai/kg diet. Mean-measured concentrations were <25.0 (<LOD, control), 555, 982, 1790, 3200, and 5290 mg ai/kg diet, respectively. No treatment-related mortality, clinical signs of toxicity, or effects on body weight or food consumption were indicated during the study. The 8-day acute dietary LC<sub>50</sub> was >5290 mg ai/kg diet, and the NOAEC was 5290 mg ai/kg diet. DPX-MAT28 Technical (aminocyclopyrachlor) would be classified as practically non-toxic to juvenile mallard duck (*Anas platyrhynchos*) in accordance with the classification system of the U.S. EPA.

This toxicity study is scientifically sound and classified as acceptable and, thus, satisfies the guideline requirement for an acute dietary toxicity study for the mallard duck.

**Results Synopsis**

Test Organism Size/Age(Mean Weight): 10-days old; 132 to 178 g

LC <sub>50</sub> : >5290 mg ai/kg diet	95% C.I.: N/A
Probit Slope: N/A	95% C.I.: N/A

NOAEC: 5290 mg ai/kg diet  
LOAEC: >5290 mg ai/kg diet

Endpoints affected: none

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical (Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

**I. MATERIALS AND METHODS**

**GUIDELINE FOLLOWED:** U.S. EPA Ecological OPPTS No. 850.2200 (1996)  
U.S. EPA Pesticide Assessment Guidelines, §71-2 (1982)  
ASTM Standard E857-87 (1987)  
OECD Guideline No. 205 (1984)

Deviations from OPPTS Guideline No. 850.2200 included:

1. The pre-test health of the population (including mortality) was not reported. Birds should not be used if >5% of the total test population die during the 72-hour period preceding the test.
2. It was unclear if a brooder temperature gradient (of approximately 22 to 38°C) was provided for the ducklings.
3. The photoperiod (16 h light:8 h dark) was slightly different than recommended (14 hr light:10 hr dark).

These deviations do not affect the scientific soundness or acceptability of the study.

**COMPLIANCE:** Signed and dated GLP, Quality Assurance, and Data Confidentiality statements were provided. This study was conducted in accordance with the GLP as published in 40 CFR Part 160 with the following exception: periodic analyses of feed and water for potential contaminants were not conducted in accordance with GLP, but were performed using a certified laboratory and standard U.S. EPA analytical methods.

**A. MATERIALS:**

**1. Test Material** DPX-MAT28 Technical (aminocyclopyrachlor)

**Description:** Solid

**Lot No./Batch No.:** Not reported; Reference No. DuPont-21466

**Purity:** 92.2%

**Stability of compound under test conditions:** The 5-day stability of DPX-MAT28 Technical was determined under actual use conditions. Recoveries averaged 95 to 102% of initial measured concentrations.

**Storage conditions of test chemicals:** Ambient conditions

**Physicochemical properties of Aminocyclopyrachlor.**

Parameter	Values	Comments
Water solubility at 20°C	Not reported	
Vapor pressure	Not reported	
UV absorption	Not reported	

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical  
(Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

Parameter	Values	Comments
pKa	Not reported	
Kow	Not reported	

(OECD recommends water solubility, stability in water and light, pKa, Pow, and vapor pressure of test compound)

**2. Test organism:**

**Species (common and scientific names):** Mallard duck (*Anas platyrhynchos*)  
(EPA recommends using either bobwhite quail or mallard duck.)

**Age at study initiation:** 10 days old  
(EPA recommends: 10-14 days old)

**Weight at study initiation (mean and range):** 132 to 178 g

**Source:** Whistling Wings, Inc., Hanover, IL

**B. STUDY DESIGN:**

**1. Experimental Conditions**

a. Range-finding Study: None reported. The dietary concentrations were established based upon known toxicity data.

b. Definitive Study:

**Table 1: Experimental Parameters**

Parameter	Details	Remarks
		Criteria
<u>Acclimation</u> Period:  Conditions: (same as test or not)  Feeding:  Health: (any mortality observed)	8 days  Same as test  Game bird ration formulated to Wildlife International Ltd.'s specifications and water from the town of Easton public water supply were offered <i>ad libitum</i> .  Not reported.	All birds were observed daily during acclimation, and any birds exhibiting abnormal behavior or physical injury were not used for the study.

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical  
(Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

Parameter	Details	Remarks
		Criteria
Pen size and construction materials	Vinyl-coated wire grid cages measuring 62 x 92 x 25.5 cm.	Birds were housed in groups of five. The floor area was approximately 1141 cm <sup>2</sup> per bird.  <i>Recommended pen size is about 35 x 100 x 24 cm</i>
Test duration	5 days with treated feed followed by 3 days with untreated feed	  <i>Recommended test duration is 5 days with treated feed and at least 3 days observation with "clean" feed.</i>
<u>Test concentrations</u> nominal:  measured:	0 (negative control), 562, 1000, 1780, 3160, and 5620 mg/kg diet  <25 (<LOD, control), 555, 982, 1790, 3200, and 5290 mg ai/kg diet, respectively	Dietary concentrations were adjusted for the purity of the test substance.  Samples were collected for analysis on Day 0. Mean-measured concentrations ranged from 94 to 101% of nominal.  <i>Five or six test concentrations should be used in a geometric scale, unless the LC<sub>50</sub> &gt; 5000 mg ai/kg diet.</i>
<u>Solvent/vehicle, if used</u> type: amount:	N/A	  <i>Recommended solvents include distilled water, corn oil, propylene glycol, 1% carboxymethylcellulose, or gum arabic. The solvent should not be more than 2%.</i>
Diet preparation and feeding	The appropriate amount of test substance was blended for approximately 1 minute with a small portion of basal diet. This mixture was then quantitatively combined with the remaining basal diet and mixed for a total of 20 minutes using a Hobart mixer.	Diets were prepared on the day of test initiation and presented to the birds.  <i>The control group should be tested with a diet containing the maximum amount of vehicle used in treated diets.</i>
Feed withholding period	None	
Stability and homogeneity of test material in the diet determined (Yes/No)	Stability – yes Homogeneity – yes	

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical  
(Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

Parameter	Details	Remarks
		Criteria
<u>Number of birds per replicate/groups</u> for negative control: for vehicle control: for treated:	5 birds/cage N/A 5 birds/cage	<i>The recommended number of birds per replicate is a minimum of ten.</i>
<u>Number of replicates/group (if used)</u> for negative control: for vehicle control: for treated:	6 replicate cages N/A 2 replicate cages/level	There were a total of 30 control birds and 10 treated birds/level, fulfilling OPPTS guidelines.
<u>Test conditions</u> temperature:  relative humidity(%):  photoperiod:	Brooder: <i>ca.</i> 29°C Room: 22.7 ± 0.8°C  Room: 33 ± 11%  16 hours light/8 hours dark	It is recommended that a temperature gradient in the pen of approximately 22 to 38°C is provided to allow young birds to seek a proper temperature.  The photoperiod slightly exceeded recommendations (14 hours light/10 hours dark); light intensity was approximately 215 lux.  <i>Recommended brooder temperature is about 35°C (95°F)</i> <i>Recommended room temperature is 22-27°C (71-81°F)</i> <i>Recommended relative humidity is 30-80%</i> <i>Recommended photoperiod is 14 hours of light/10 hours dark.</i>
Reference chemical, if used	N/A	

**2. Observations:**

**Table 2: Observations**

Parameters	Details	Remarks
Parameters measured (mortality/body weight/mean feed consumption/others)	- Mortality - Clinical signs of toxicity - Body weight - Food consumption	

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical (Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

Parameters	Details	Remarks
Indicate the stability and homogeneity of test chemical in the diet	<p><u>Stability:</u> Stability was assessed at all levels after 5 days under actual use conditions. Recoveries averaged 95 to 102% of initial measured concentrations.</p> <p><u>Homogeneity:</u> Homogeneity was assessed by collecting samples from the top, middle, and bottom left and right areas (six samples) from treated feed prepared at 562 (low) and 5620 (high) mg/kg diet. Coefficients of variation were 1.68 and 2.11%, respectively.</p>	
Indicate if the test material was regurgitated	No regurgitation was indicated.	
Treatments on which necropsies were performed	None performed	As no mortalities were observed, gross necropsies were not necessary.
Observation intervals	Birds were observed at least twice daily for mortality and clinical signs of toxicity. Body weights were determined on Days 0, 5, and 8. Feed consumption was calculated each day during the exposure period and once during the recovery period (Days 6 to 8).	
Were raw data included?	Raw body weight and food consumption data were not provided.	

## II. RESULTS AND DISCUSSION:

### A. MORTALITY:

No mortality was observed in the control or any treatment group. The 8-day acute dietary LC<sub>50</sub> was >5620 mg ai/kg diet. The NOAEC for mortality was 5620 mg ai/kg diet.

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical (Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

**Table 3: Effect of DPX-MAT28 Technical (Aminocyclopyrachlor) on Mortality of Mallard Duck.**

Treatment, mg ai/kg diet Mean-measured (and nominal) conc.		No. of birds per treatment	Cumulative mortality				
			day 1	day 2	day 3	day 4	day 8
Control		30	0	0	0	0	0
555 (562)		10	0	0	0	0	0
982 (1000)		10	0	0	0	0	0
1790 (1780)		10	0	0	0	0	0
3200 (3160)		10	0	0	0	0	0
5290 (5620)		10	0	0	0	0	0
NOAEC		5620 mg ai/kg diet (nominal)					
LC <sub>50</sub>		>5620 mg ai/kg diet (nominal)					
Reference chemical	mortality	N/A					
	LC <sub>50</sub>	N/A					
	NOAEC	N/A					

**B. SUB-LETHAL TOXICITY ENDPOINTS:**

There were no overt signs of toxicity at any of the concentrations tested. With one exception, all birds from all treatment and control groups were normal in appearance and behavior throughout the test. One bird in the 1000 mg ai/kg diet group was observed limping in the morning of Day 1 due to having its leg caught in the cage. The bird appeared normal during the afternoon observation and for the remainder of the study. The NOAEC for clinical signs of toxicity was 5620 mg ai/kg diet.

Visual inspection of the data indicated no apparent treatment-related effects upon body weight or food consumption at any treatment level. The NOAEC values for body weight and feed consumption data were 5620 mg ai/kg diet.

Using body weight and food consumption data, daily dietary doses were estimated to be 208, 439, 779, 1530, and 2423 mg ai/kg/day for the 562, 1000, 1780, 3160, and 5620 mg ai/kg diet levels, respectively.



**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical (Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

**Table 4: Sublethal Effect of DPX-MAT28 Technical (Aminocyclopyrachlor) on Mallard Duck.**

Treatment, mg ai/kg diet Mean-measured (and nominal) conc.		Observation									
		Body weight change, g			Food consumption, g/bird/day						
		Days									
		0-5	5-8	0-8	0-1	1-2	2-3	3-4	4-5	0-5	6-8
Control		158	91	248	97	90	102	114	137	108	150
555 (562)		161	88	249	65	83	86	91	106	86	126
982 (1000)		150	99	249	74	90	98	113	131	101	162
1790 (1780)		159	101	260	74	90	92	112	127	99	149
3200 (3160)		152	96	247	84	100	105	113	149	110	148
5290 (5620)		170	89	258	75	93	97	117	132	103	153
NOAEC		5620 mg ai/kg diet (nominal)			5620 mg ai/kg diet (nominal)						
EC <sub>50</sub>		>5620 mg ai/kg diet (nominal)			>5620 mg ai/kg diet (nominal)						
Reference chemical	NOAEC	N/A			N/A						
	EC <sub>50</sub>	N/A			N/A						

**C. REPORTED STATISTICS:**

As no mortalities were observed, the 8-day LC<sub>50</sub> was visually determined to be greater than the highest concentration level. No statistical analyses were applied to separate mean responses among treatment groups for the endpoints of food consumption and body weight.

**D. VERIFICATION OF STATISTICAL RESULTS:**

Statistical Method: Statistical analyses were not required to verify the results of this study. There was no mortality and body weight and food consumption were clearly not affected by treatment in any dose-dependent manner.

LC<sub>50</sub>: >5290 mg ai/kg diet      95% C.I.: N/A

Probit Slope: N/A      95% C.I.: N/A

NOAEC: 5290 mg ai/kg diet

Endpoints affected: none

**E. STUDY DEFICIENCIES:**

There were no significant deviations from U.S. EPA OPPTS Guideline No. 850.2200 affecting the scientific soundness or acceptability of this study.

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical  
(Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

**F. REVIEWER'S COMMENTS:**

The reviewer's conclusions agreed with those of the study authors. Results were provided in terms of mean-measured concentrations in the Executive Summary and Conclusions sections of the DER.

Portions of treated feed (10.0 g) were extracted with 100 mL of methanol:water (80:20, v:v) by sonicating for 15 minutes followed by shaking at *ca.* 250 rpm for *ca.* 60 minutes. Aliquots were centrifuged at *ca.* 1500 rpm for *ca.* 10 minutes, diluted as needed with methanol:water (80:20, v:v), and analyzed for aminocyclopyrachlor using HPLC with UV (240 nm) detection. The analytical LOD and LOQ were 25.0 and 50.0 mg ai/kg diet, respectively.

Avian diet was fortified at 50.0 or 6000 mg ai/kg and analyzed concurrently with the Day 0 and 5 samples. Mean recoveries were 93% for both days. Test sample results were not corrected for the mean procedural recoveries from each sample set.

Experiment study dates were March 29 to April 6, 2007.

**G. CONCLUSIONS:**

This study is scientifically sound and classified as acceptable. No treatment-related mortality, clinical signs of toxicity, or effects on body weight or feed consumption were observed at any test level following a 5-day acute dietary administration of DPX-MAT28 Technical (aminocyclopyrachlor) at up to the limit concentration, followed by a 3-day recovery period. The  $LC_{50}$  was >5290 mg ai/kg bw (>limit concentration), and the NOAEC was 5290 mg ai/kg bw.

$LC_{50}$ : >5290 mg ai/kg diet	95% C.I.: N/A
Probit Slope: N/A	95% C.I.: N/A

NOAEC: 5290 mg ai/kg diet  
LOAEC: >5290 mg ai/kg diet

Endpoints affected: none

**Data Evaluation Report on the Acute Dietary Toxicity of DPX-MAT28 Technical  
(Aminocyclopyrachlor) to Mallard Duck (*Anas platyrhynchos*)**

PMRA Submission Number {.....}

EPA MRID Number 47560119

---

**III. REFERENCES:**

U.S. Environmental Protection Agency. 1996. Series 850 – Ecological Effects Test Guidelines (*draft*), OPPTS Number 850.2200: *Avian Dietary Toxicity Test*.

U.S. Environmental Protection Agency. 1982. *Pesticide Assessment Guidelines, FIFRA Subdivision E, Hazard Evaluation: Wildlife and Aquatic Organisms*, Subsection 71-2, Environmental Protection Agency, Office of Pesticide Programs, Washington, D.C.

American Society for Testing and Materials. 1987. Standard Practice for Conducting Subacute Dietary Toxicity Tests with Avian Species. ASTM Standard E857-87. Annual Book of ASTM Standards, Vol. 11.04. Philadelphia, PA.

Organization for Economic cooperation and Development. 1984. *Avian Dietary Toxicity Test*. OECD Guideline for Testing of Chemicals. Guideline 205. Paris.

National Research Council. 1996. *Guide for the Care and Use of Laboratory Animals*. Washington, D.C. National Academy Press. 125 pp.

Finney, D.J. 1971. *Statistical Methods in Biological Assay*, Second Edition, Griffin Press, London.

Thompson, W.R. 1947. *Bacteriological Reviews*. Vol. II, No. 2 (June): 115-145.

Stephan, C.E. 1977. Methods for Calculating an LC<sub>50</sub>. Pages 65-84 in *Aquatic Toxicology and Hazard Evaluations*, American Society for Testing and Materials. Pub. No. STP 634. Philadelphia, PA.

Stephan, C.E. 1978. U.S. EPA, Environmental Research Laboratory, Duluth, Minnesota. Personal Communication.

